

---

*Note:* You will be given time to solve the exercises during the tutorial.

### Exercise 1.1: Estimate $\pi$ with Monte Carlo

- Estimate  $\pi$  by “shooting” (i.e., drawing random numbers)  $N$  times uniformly on a square and counting the number of points hitting a disc target: the ratio of hits to  $N$  should correspond to the ratio of the areas of the target to the area you shoot on.
- Estimate the variance of the error for given  $N$  by repeating this a few times.
- Plot the variance of the error versus  $N$  on a log-log scale. What is the scaling of the error?

### Exercise 1.2: Metropolis algorithm for the 2D Ising model

Download the script `metropolis.py` from the homepage, which implements the Metropolis algorithm for the classical 2D Ising model  $H = -J \sum_{\langle i,j \rangle} \sigma_i \sigma_j$  with  $J \equiv 1$ . The 2D Ising model has a critical point at  $T_c = 2J / \ln(1 + \sqrt{2}) \approx 2.269$ .

- What is the script plotting?
- What are “typical” configurations at temperatures  $T \gg T_c$ ,  $T \approx T_c$  and  $T \ll T_c$ ?
- Plot the energy  $E$  and specific heat  $C_V$  versus temperature  $T$  for different system sizes  $L$ .
- Adjust the script to measure the magnetization  $M = \frac{1}{L^2} \sum_{i,j} \sigma_{i,j}$ . Plot how  $M$  changes with simulation time (=the number of updates performed) for  $T > T_c$ ,  $T \approx T_c$  and  $T < T_c$ . Which time scales can you recognize? In which cases do you still get the correct expectation value  $\langle M \rangle = 0$ ? Plot  $\langle |M| \rangle$  (i.e. taking the absolute value of  $M$  before averaging) versus  $T$  to see the transition.
- Include a magnetic field  $h$  coupling to the spins with a term  $H' = -h \sum_i \sigma_i$ . Plot  $\langle M \rangle$  versus  $T$ .

Bonus Some further ideas for playing around:

- Instead of restarting from a random state for each new  $\beta$ , re-use the last state of the previous simulation. You should still perform sweeps without measurements for the thermalization. Is it better to start with large  $\beta$  or small  $\beta$ ?
- Change the lattice.
- Optimize the code.
- ...