Clarical Field theory: spin-1 fields

> QFT WS 2025 Leurs o Tound

As for as we know, matter is made by spin-1 elementary components => electrons, quaks, protons... let us there for spen to how we can build a field theory for spen to (clonical) fields We have seen that fundamental building blacks from the point of view of Poincre are Weyl spinoss 1 test with (10) LEFT-HANDED 4/2 First, we need a KINETIC TERM which is breutz imuariout => must mustre 2µ oud so we med a 4-vector to get a SCALAR ? We hove seen that 4 5" 4" behoves indeed. es 4- vector ander Lorent t (see Lecture 3) Sim losly $\frac{1}{\sqrt{R}} \sigma^{M} + \frac{1}{\sqrt{R}} s do u - vector$ and remember $\sigma^{M} = (4L, \vec{\sigma})$ $\vec{\sigma}^{M} = (4L, -\vec{\sigma})$

=> con we build something else? 7 7 2 IS NOT LORENTZ INVARIANT we would need 4the or the 4L We can there fre wild simplest corent & scalor or [4] = 3 1 2 K = 1 4 5 m 2 14 L mon - dimension Where the i is needed to get Jd'x LK ER (14t 5 m 2, 41) = -i [4t 5 m 2, 41]* = -i [], 4,] t (5") + 4. = -i るルレナーナーナーナーナーコルナー where I used (5m) = 5m hermiticity

Z

If we coulder this on the complete of them Equations of motion become 14 - 9 3x 46 + 146 北: => 5 4 24 = 0 Wyl Equation FIRST ONDER! let's now multiply this by ord w theu Using 50 54 2 10 = 1 [50 54 + 540] 2 20 = 1100, 5432 000 $\begin{cases}
31 & \text{for } \mu = V = 0 \\
11, -\sigma^{i} = -2\sigma^{i} & \text{for } \mu = i, V = 0 \\
16', 113 = +3\sigma^{i} & \text{fr } \mu = 0, V = 1 \\
-(\sigma^{i}, \sigma^{2}) = -2\delta^{i} & \text{for } \mu = i, V = 1
\end{cases}$

.

then
$$\frac{1}{2}$$
 for, $\frac{1}{6}$ $\frac{1}{9}$ $\frac{1}$

$$\frac{2}{2}(p) = u_{L}(p) e^{-ip \times} = p^{2} = 0$$
Experience
but we also have 1st order equation, which is
$$\frac{1}{2}(p) = u_{L}(p) e^{-ip \times} = 0$$

$$\frac{1}{2}(p) = u_{L}(p) e^{-ip \times} = 0$$

 $\Rightarrow \overline{\sigma}^{\mu} P_{\mu} U_{L} = 0 \quad (E_{\rho} + \overrightarrow{\sigma} \cdot \overrightarrow{P}) U_{L} = 0$ because $\overline{\sigma}^{\prime} = (U_{\rho} - \overrightarrow{\sigma})!$

The spin
$$\frac{1}{2}$$
 with $\frac{1}{2}$ remainder now that for spin $\frac{1}{2}$ field the augular momentum is given by $\vec{J} = \frac{\vec{r}}{2}$
 $\vec{p} \cdot \vec{J} \cdot U_L = -\frac{1}{2} \cdot U_L$ with $\vec{p} = \frac{\vec{p}}{|\vec{p}|}$

which means that $\frac{1}{4}$ has projection of spin along direction of momentum = $-\frac{1}{2}$. Thus HELICITY = $-\frac{1}{2}$, which is

what we expect from Little group monten spin ?!

For now this is a statement about the Weyl feld which belongs to (½,0) irres of LORENTZ

=> After Quartitation +'00 became don't PARTICLES!

We can compose the energy -momentum tower 8m = 32 34 + 34+ 3k - gnv 2 become ou E.M X = 0 5 m 2, 4 = 0 = かんちゅうなし hich that $\theta^{00} = i \chi^{\dagger} \partial^{3} \psi_{L} \sim 7$ HAMILTONIAN Also, ano this is a complex FIELD, we can do earth U(1) transformation 2/2 > e18 +2

And that the conserved change is

Q = Sd² × 4t 4t this care aprim be interpreted as electric dage!

(see Later!)

Write also that we can stort from EQUIVALENT of the of the office of the

$$= \frac{i}{2} \left[\frac{1}{4} + \frac{1}{5} + \frac{3}{4} + \frac$$

Conserved current looks different, but charge one the source! EXERCISE Now of course this theory is NOT parity invoicent more P sends 42 -> 2/R for let's frot construct the equivolent theory.

for a RIGHT-HANDED WEYL SPINOR LR = 1 YR 5 m 2 YR where similarly E.H OM Dute = 0 => II tr = 0 multiply by 5 2 tag su etmograma to paticu won tod $y_e = u_R e^{-ip \times x}$ then $p^2 = 0$; $f_p = |\vec{p}|$ -10 pm UR =0 => (Ep - 5.7) UR =0 $\vec{J} \cdot \hat{\vec{p}} = + \frac{1}{2} u_R$ helichy $+\frac{1}{2}$.

DIRAC FIELDS

We would like to construct now a theory with both the diff => PARITY INVARIANCE ?

now I can consider of the diff the little diff the last they are noticed man forms

prendo - scolar exclude for P-EVEN 9 =) We focus on pority EVEN Theory (Draw thery) Knetic port is PARITY EVEN becouse P: 42 > 42 P: 42 > 42 P: 24 -> (30,-2i) => P: 2,50 = 2,50 Fix Qo = 0; QE = - m, os we see now, mon tem As usual, we can work out Eulen-Lagrange Eq.s

As before, take first Eq and multiply (it dv)

 $= - \frac{1}{2} \{ (\nabla^{2} - \nabla^{2} - \nabla^{2}$

10 we get - 172 = m24L => (1+m2)42 =0 massive klein-gordon Eq! mon porometer for 4, 4, and sho $(U+m^2) \forall R=0$ We can now reallest everything as terms of one simple DRAC SANDR $\gamma_{\bar{0}}^{-} \left(\begin{array}{c} \gamma_{L} \\ \gamma_{R} \end{array} \right) = \left(\begin{array}{c} \gamma_{L} \\ \gamma_{R} \end{array} \right) \oplus \left(\begin{array}{c} \gamma_{L} \\ \gamma_{L} \end{array} \right) \oplus \left(\begin{array}{c} \gamma_{L} \\ \gamma_{R} \end{array} \right) = 0$ diamised this is a special representation for Drac's Spinors colled CHIRAL REPRESENTATION , we will see more representations later on. lu this representation we can DEFINE Diac's & matical $\chi_{\text{M}} = \begin{pmatrix} Q_{\text{M}} & Q_{\text{M}} \\ Q_{\text{M}} & Q_{\text{M}} \end{pmatrix} \Rightarrow \chi_{\text{M}} = \begin{pmatrix} Q_{\text{M}} & Q_{\text{M}} \\ Q_{\text{M}} & Q_{\text{M}} \end{pmatrix} \xrightarrow{\text{NAME/X}} 11$

CUFFORD 1 8h 8 = 2 gm 11 4x4 ruch that ALCIEBRA Droc Spinor space => 8th 4x4 motrices, smallest possible DIMENSION!

(more LATER) With this notation the two "Weyl" massive eps become (i 3 / 1/m) 2+0 = 96 $(i \not > -m) \not \downarrow_{D} = 0$ where $\not > = \xi^{\mu} \not >_{\mu}$ DIRAC EQUATION We can also derive this equation directly from a L 24 (i) -m) 4 Lo= 40 (17-m) 40 Drop "D" for simplicity And we defined DIRAC ADJOINT 7= 48

 $\Rightarrow \begin{cases} 0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \end{cases}$ is necessary to have right Kinetic d'mon terms in terms of 42 d 4/2!

XI matrices are a representation of <u>Cect</u> 3) Interestingly consider Zw= 4[7, 7,]

You can prove that [5m, 2po] = i (gre 2no - gre 2no - gro 2ne + gro 2re) (Exelcise)

~> Clifford olgebra "contains" a representation of

remember we wrote for a borentz transformed a (see Locture 3) $\Lambda = e^{-\frac{i}{2}\omega_{\mu\nu}J^{\mu\nu}}$

then we can generate a REPRESENTATION of Lorents on DIRAC-SPINOR SPACE

$$\Lambda \beta = \beta' = e^{-\frac{i}{2}\omega_{\mu\nu}} \sum_{\mu\nu}^{\mu\nu}$$

$$\int generators on Spinore$$
SPACE

and one can see explicitly that

 $\Theta^{i} = \frac{1}{2} \, \epsilon^{ijk} \, \omega^{ijk} \quad ; \quad \eta^{i} = \omega^{i0}$

$$\Rightarrow$$
 of course, this is a REDUCISLE representation time Dirac = $\left(\frac{1}{2}, 0\right) \oplus \left(0, \frac{1}{2}\right)$

Now back to Dirac Equation of Dirac Spinors

We define a 5th y matrix
$$75$$
 $75 = 1709^{4}y^{2}y^{3} = (-12x^{2})$ in chiral representation

with 57^{μ} , $763 = 0$ $=$ representation.

With this we see
$$P_L = \frac{1-15}{2} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$P_R = \frac{1+15}{2} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

Project out left & hight components of spinors

=> Pr. Pr = Pr; Pr Pr = Pr; Pr Pr = Pr Pre = Pr Pre = Pre = 0

projector operators

Pry= (4) Pry= (4)

Pry= (4)

Now, early we have guessed the form of LD using usual requirements? => Great invariouse etc Stort noticing that & is axa matrix . a general 4×4 matrix hos 16 (couplex) entres . We can build 16 independent 4×4 matrices {1, 8", 85, 8" 85, 5" } 5 MV = - 5 VM 10 matica 6 independent oves => Any uxu H = lines combination of these 16

this allows us to clonify all possible terms our a Lapon gan modering Direct Sproks

We have seen \(\frac{1}{4} = \frac{1}{4} + \

exclude for pseudo Scalar under full
P-invoiant Loventz group (including P)
theory! Loventz group (including P)
then we had: If x " 4" = 4", 5" 4; + 4" 5" 4;

then we find: If y'may = 4t 5 myz + 4to myz
both 6-vectors

this must
transform as a 6 vector = 1" "VECTOR"
WELLEUT"

notice that under P 42 4242
which lauplies 10 > 10; 10 > - 10 4-vector!

· on the other houd: 7 x5 xmy = 4t 5 m21 - 7komyk = JA AMAL-VECTOR

WELENT is a pseudo vector becouse Ja - Ja domles

Ja - Ja domles

PARITY! => exclude it for PARITY INVARIANT THERY · findly 4 2 mg - 10 a 6-tensor Sangu not us => From this we see that we end up

FREE DIRAC EQUATION b=du=0; d Cm = i du EXTRA PIECE 7 5 my requires something to contract two indiceo: "epi" which is ANTIONNETRIC -> We can use FMV => epr= e Fmv non-minimal aupling HAGNETIC PIPOLE WTELACTION agoin "suppressed" > LATER!

Folder os fows on our FREE DRAC LAGRANGIAN 7 (i yu 2 m) 7 We have seen that you generale CLIFFORD ALGEBRA oud we found one specific representation $\chi^{M} = \begin{pmatrix} 0 & 0^{M} \\ \overline{6}^{M} & 0 \end{pmatrix} \qquad \chi^{5} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \qquad \text{REPR.}$ in which $y = \begin{pmatrix} y_L \\ y_R \end{pmatrix}$ this is not the only one: onsider 4 = 04 U unitary matex

20 = 21 Upo Ut U [i χη θμ -m] Ut 4 1

= [2+ xp]' [i θμ χμ - m] 24 1

= 7' [i θμ χμ - m] 24 1

with χ'' in new representation $\chi'' = U \chi'' U^{\dagger}$ Indeed this transformation does not drouge cufford MC.

[$\chi'' \chi'' = \chi'' \chi'' = \chi'' \chi'' = U [\chi'' \chi'' + \chi'' \chi'' = U [\chi'' + \chi'' \chi'' = U [\chi'' + \chi'' \chi'']]]$

go indeed y" still represent some elgebra. Diroc
Equation s invariant: La des invariant!

Let us cansider a specific choice for $U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$ $Y' = \frac{1}{\sqrt{2}} \begin{pmatrix} 4 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 4 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix}$ The standard representation representation

This case $\chi^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ $\chi^{*} = \begin{pmatrix} 0 & 0^{-1} \\ -0^{-1} & 0 \end{pmatrix}$

$$8^{5} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

In this representation, 4, the ore more HIDDEN,

no it's less convenient in that ENERGY LIMIT

There are EICENSTATES of helicity for M=0

They are EICENSTATES of helicity for M=0. For LOW ENERGY, m =0 and helicity is not one eigenstate, in this case STANDARD REPR.

Is more useful

What short solution of the free Dirac Equation?

L & R satisfy Klen-gordon Eq We have seen that multiply by - if => (i3-m)4=0 + 3, 2, 8 4 + m i 24 = 0 17 + m27 = 0 (1) +m2) += 0 10 agains general volution con le decomposed in Forcier modes

The again period relation can be decomposed in Forcier mode
$$\frac{2}{\sqrt{1+x}} = \frac{2}{5} \int \frac{d^3\vec{p}}{(2\pi)^3} \left(2E_p\right) \left[2s(p) \cdot \frac{1}{9}(p) \cdot e^{-ip \cdot x}\right]$$

$$+ \sqrt{1+x} \cdot \left[2s(p) \cdot \frac{1}{9}(p) \cdot e^{-ip \cdot x}\right]$$

Us(p) Vs(p) (6-Din) SPINORS S=1,2 SPINS!

Us(p), Vs(p) & Opis, Lipis ore différent becouse T CONCEX FIELD! => I could door by L > 4, V but convenient for QUANTIZATION ?

$$(i\beta-m)\psi=0 \qquad (y+m) \, v_s(p)$$

$$u(p) = \begin{pmatrix} u_L \\ u_P \end{pmatrix}$$
 and assume $m \neq 0$
then we go to rest frame $p^{\mu} = (m, \vec{o})$

1]
$$m[y^{0}-1]u_{s}(p)=0$$

2] $m[y^{0}+1]v_{s}(p)=0$

$$\Rightarrow 1] \quad u_s = u_s \qquad 2] \quad v_s^R = -v_s^L \quad \forall$$

Define BASIS
$$\begin{cases} S_{1} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ S_{2} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \end{cases}$$

UL=UR=Vm &

oud vormolize

$$U_{S}(p) = \left[\left[\frac{1-\sigma_{3}}{2} \right] + \sqrt{E-P_{2}} \left(\frac{1+\sigma_{3}}{2} \right) \right] \xi_{S}$$

$$\left[\left[\frac{1+\sigma_{3}}{2} \right] + \sqrt{E-P_{2}} \left(\frac{1-\sigma_{3}}{2} \right) \right] \xi_{S}$$

$$\sigma_{3} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \qquad \frac{1-03}{2} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \qquad \frac{1+03}{2} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \qquad 24$$

Consider the high-energy limit
$$P_{\mu} \rightarrow (E, 0, 0, E)$$
:

 $V_{2E} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$
 $V_{3s} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$
 $V_{2E} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$
 $V_{2E} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$

$$U_1(p) \rightarrow \sqrt{2E} \begin{pmatrix} 0 \\ S_1 \end{pmatrix}$$
 $U_2(p) \rightarrow \sqrt{2E} \begin{pmatrix} S_2 \\ 0 \end{pmatrix}$

only RIGHT HANDED only LEFT HANDED

note that
$$S_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$
 SAN UP; $S_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ SPIN DOWN

Similarly (=)
$$V_{5}(p)$$

$$\left(\left[V_{E+p_{2}} \left(\frac{1-63}{2} \right) + V_{E-p_{2}} \left(\frac{1+63}{2} \right) \right] \eta_{5} \right) \quad \eta_{4} = \binom{1}{0}$$

$$- \left[V_{E+p_{2}} \left(\frac{1+63}{2} \right) + V_{E-p_{2}} \left(\frac{1-63}{2} \right) \right] \eta_{5} \right) \quad \eta_{2} = \binom{0}{1}$$

$$\gamma_{2} = \binom{0}{1}$$

$$\gamma_{3} = \binom{0}{1}$$

$$\gamma_{4} = \binom{0}{1}$$

$$\gamma_{5} = \binom{0}{1}$$

$$\gamma_{5} = \binom{0}{1}$$

Different Sign on USCP)!

8, 5, l

and we can define $\overline{u}_s = u_s^t y^s$ $\overline{v}_s = v_s^t y^s$ => \overline{v}_s rac Adjoint Spinors

the normalitation us chose imples

$$\overline{\mathcal{U}}_{r}(p) \, \mathcal{U}_{s}(p) = 2m \, \delta_{rs} \, \overline{\mathcal{V}}_{r}(p) \, \mathcal{V}_{s}(p) = -2m \, \delta_{rs}$$

$$\overline{\mathcal{U}}_{r}(p) \, \mathcal{U}_{s}(p) = 2E_{p} \, \delta_{rs} \, \overline{\mathcal{V}}_{r}(p) \, \mathcal{V}_{s}(p) = 2E_{p} \, \delta_{rs}$$

+ Dirac ORTHOGONALITY $\sqrt{u_r(p)} v_s(p) = 0$ $\sqrt{v_r(p)} u_s(p) = 0$

note that we can also rewrite spinors as
$$u_{s}(p) = \begin{pmatrix} \sqrt{p \cdot \sigma} & \xi_{s} \\ \sqrt{p \cdot \overline{\sigma}} & \xi_{s} \end{pmatrix} \qquad v_{s}(p) = \begin{pmatrix} \sqrt{p \cdot \overline{\sigma}} & \eta_{s} \\ -\sqrt{p \cdot \overline{\sigma}} & \eta_{s} \end{pmatrix}$$

=>VHATRIX ruples

PSITTUE ROOT OF EACH EIGENVALUE

this form is Valid due for p^{μ} not slong z-direction => remember $(p \sigma)(p \overline{\sigma}) = p^{\mu}p^{\nu} \sigma_{\mu} \overline{\sigma}_{\nu} = p^2 = m^2$ SPIN SUMS

For laken, it will be useful to be able to SUM over all polonizations of postules involved

$$\frac{2}{s=1,2} u_{s(p)} \overline{u}_{s(p)} = \frac{2}{s} \left(\begin{array}{c} \sqrt{p \cdot \sigma} & S_{s} \\ \sqrt{p \cdot \sigma} & S_{s} \end{array} \right) \left(\begin{array}{c} S_{s} \sqrt{p \cdot \sigma} \\ S_{s} \sqrt{p \cdot \sigma} \end{array} \right) \\
= \frac{2}{s} \left[\begin{array}{c} m & S_{s} S_{s} \\ p \cdot \overline{\sigma} & S_{s} S_{s} \end{array} \right] = \left[\begin{array}{c} m & p \cdot \overline{\sigma} \\ p \cdot \overline{\sigma} & m \end{array} \right] \\
= \frac{2}{s} \left[\begin{array}{c} m & p \cdot \overline{\sigma} \\ p \cdot \overline{\sigma} & S_{s} S_{s} \end{array} \right] = \left[\begin{array}{c} m & p \cdot \overline{\sigma} \\ p \cdot \overline{\sigma} & m \end{array} \right]$$

becomes $\frac{3}{3}$ $\frac{5}{3}$ $\frac{5}{3$

Similarly we get

The DIRAC LAGRANGIAN

has a last comment, the DIRAC LAGRANGIAN

has an extra set of Global Symmetries

let us go back to CHRAL BASIS

imagine now to generalit U(1) transform. I complex scalor field as follows

this is dearly a symmetry of LD if m = 0 note now two special cones

· The = De = a (the) = eid (the) or Dience

The server

y - etd 7

the Rs is invariant under this transformation even if m to !

· Da=-OL=B (M) - (eiby) = eibys (th)

because $1/5 = \begin{pmatrix} -1 & 0 \\ 0 & 4 \end{pmatrix}$ exercise: prove that they one

symmetric using 20 in terms of 24 Symmetry only if m = 0

In fet myy = m(y+ yo 24) = m[e+85 y] xo eif85 y

= my e + 2 i 685 y not invoicet!

the Noether currents associated to the two

1"= 78"4 for U(1) vector eight
1"= 48"4 for U(1) axial eight

Symmetries one

The Tong of the O(1) exial eight of this is alled a CHIRAL TRANSFORMATION & WE say that Symmetric

Note the 21 of 1 = 21 m + x5 = m breaks

Exercise 7 30

Haking UU), a LOCAL SYNHETRY

When discussing the Scalar (complex) field, we have seen that to promote GLOBAK U(1) to a LOCAL symmetry we need to introduce a new spin 1 vector field Am I connection I which defines a covaciant description

dμφ → Dμφ = (dμ + iq Aμ) φ

charge

electromagnetic

field!

If you go back to our derivations, you will see that nothing depends on consumer properties of the field $\phi \Rightarrow$ we can repeat the very some argument for if

of we would muariouce under LOGAL ULI) 4 -> e id(x) 4 We need to CHANGE Du > Du = Su +igA M Lo > 7[~8"D,-m]4 = 7 [10 m 2 m - 4 - 9 A m 7 m 7 minimal coupling of Ap to of through vector ament fr

Q= | dx 78°4 CIRRENT REMAINS CONSERVED es thermy still hos-GLOBAL U(1) V will become electic charge often QUANTIZATION!

if we odd to this I the knetic term for Ap we get LOED (QUANTUM) ELECTRODYNAMICS Hay to betthough ten such sw LOED = 7 LID-m J4 - 4 FNV FNV So we see again that by making global Symmetries LOCAL for SCALAR & SPINOR FIELDS (HATTER!) we notrolly generate SPIN 1 felds In the next lecture we will see what we can say chamically don't these theories ...